

<p>(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/705, C12N 1/21, 15/70, G01N 33/53, 33/68, A61K 38/17, C07K 17/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/32767</p> <p>(43) International Publication Date: 8 June 2000 (08.06.00)</p>
<p>(21) International Application Number: PCT/EP99/09440</p> <p>(22) International Filing Date: 3 December 1999 (03.12.99)</p> <p>(30) Priority Data: 98122969.3 3 December 1998 (03.12.98) EP</p> <p>(71) Applicant (for all designated States except US): MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Hofgartenstrasse 8, D-80539 München (DE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SONDERMANN, Peter [DE/DE]; Pfeufferstrasse 97, D-81373 München (DE). HUBER, Robert [DE/DE]; Schlesierstrasse 13, D-82110 Germering (DE). JAKOB, Uwe [DE/DE]; Guldeinstrasse 42, D-80339 München (DE).</p> <p>(74) Agents: WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: RECOMBINANT SOLUBLE Fc RECEPTORS</p>		
<p>(57) Abstract</p> <p>Recombinant soluble Fc receptors according to the present invention are characterized by the absence of transmembrane domains, signal peptides and glycosylation. Such Fc receptors can easily be obtained by expressing respective nucleic acids in prokaryotic host cells and renaturation of the obtained inclusion bodies, which procedure leads to a very homogenous and pure product. The products can be used for diagnostic as well as pharmaceutical applications and also for the generation of crystal structure data. Such crystal structure data can be used for the modelling of artificial molecules. A further embodiment comprises coupling the Fc receptors according to the invention to solid materials like chromatography materials that can be used to separate and/or enrich antibodies.</p>		